

ARTICLE

Putting seedlings on the map: Trade-offs in demographic rates between ontogenetic size classes in five tropical forests

Stephan Kambach^{1,2}  | Helge Bruelheide^{1,2} | Liza S. Comita^{3,4}  |
 Richard Condit^{5,6}  | S. Joseph Wright⁴  | Salomón Aguilar⁴ |
 Chia-Hao Chang-Yang⁷ | Yu-Yun Chen⁸ | Nancy C. Garwood⁹ |
 Stephen P. Hubbell^{4,10} | Pei-Jen Luo⁷ | Margaret R. Metz¹¹  |
 Musalmah Bt. Nasardin¹² | Rolando Pérez⁴ | Simon A. Queenborough³  |
 I-Fang Sun¹³ | Nathan G. Swenson¹⁴ | Jill Thompson¹⁵ | María Uriarte¹⁶ |
 Renato Valencia¹⁷ | Tze Leong Yao¹² | Jess K. Zimmerman¹⁸ | Nadja Rüger^{2,4,19}

¹Institute of Biology, Department of Geobotany and Botanical Garden, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany

²German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany

³School of the Environment, Yale University, New Haven, Connecticut, USA

⁴Smithsonian Tropical Research Institute, Panama, Ancón, Panama

⁵Field Museum of Natural History, Chicago, Illinois, USA

⁶Morton Arboretum, Lisle, Illinois, USA

⁷Department of Biological Sciences, National Sun Yat-sen University, Kaohsiung, Taiwan

⁸Department of Natural Resources and Environmental Studies, National Dong Hwa University, Hualien, Taiwan

⁹School of Biological Sciences, Southern Illinois University Carbondale, Carbondale, Illinois, USA

¹⁰Department of Ecology and Evolutionary Biology, University of California, Los Angeles, Los Angeles, California, USA

¹¹Department of Biology, Lewis & Clark College, Portland, Oregon, USA

¹²Forest Research Institute Malaysia (FRIM), Kepong, Selangor, Malaysia

¹³Center for Interdisciplinary Research on Ecology and Sustainability (CIRES), National Dong Hwa University, Hualien, Taiwan

¹⁴Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana, USA

¹⁵UK Centre for Ecology & Hydrology, Penicuik, UK

¹⁶Department of Ecology, Evolution & Environmental Biology, Columbia University, New York, New York, USA

¹⁷Escuela de Ciencias Biológicas, Pontificia Universidad Católica del Ecuador, Aptado, Quito, Ecuador

¹⁸Department of Environmental Sciences, Universidad de Puerto Rico, San Juan, Puerto Rico

¹⁹Department of Economics, University of Leipzig, Leipzig, Germany

Correspondence

Stephan Kambach

Email: stephan.kambach@gmail.com

Abstract

All species must partition resources among the processes that underly growth, survival, and reproduction. The resulting demographic trade-offs constrain the

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Author(s). *Ecology* published by Wiley Periodicals LLC on behalf of The Ecological Society of America.

Funding information

Conservation, Food and Health Foundation (CFHF); National Geographic Society (NGS); German Research Foundation; German Centre for Integrative Biodiversity Research; Forest Global Earth Observatory; DFG, Grant/Award Numbers: BR 1698/21-1, HI 1538/16-1; Smithsonian Tropical Research Institute; Swiss National Science Foundation, Grant/Award Number: 193907; 2019-2020 BiodivERsA joint call for research proposals, under the BiodivClim ERA-Net COFUND program; Technology Agency of the Czech Republic, Grant/Award Number: SS70010002; Agence Nationale de la Recherche, Grant/Award Number: ANR-20-EBI5-0001-05; Japan Society for the Promotion of Science (JSPS); Smithsonian Tropical Research Institute (STRI); National Science Foundation (NSF), Grant/Award Numbers: BSR-8811902, DEB 9411973, DEB 0080538, DEB 0218039, DEB 0620910, DEB 0963447, DEB-129764; NSF LTREB, Grant/Award Numbers: DEB 0075102, DEB 0823728, DEB 0640386, DEB 1242622, DEB 1464389; Taiwan Forestry Bureau; Taiwan Forestry Research Institute; Ministry of Science and Technology of Taiwan; University of Puerto Rico; International Institute of Tropical Forestry, USDA Forest Service; Mellon foundation; Negeri Sembilan State Forestry Department; Pontifical Catholic University of Ecuador, Grant/Award Numbers: L13251, M13373

Handling Editor: Joseph B. Yavitt

range of viable life-history strategies and are hypothesized to promote local coexistence. Tropical forests pose ideal systems to study demographic trade-offs as they have a high diversity of coexisting tree species whose life-history strategies tend to align along two orthogonal axes of variation: a growth–survival trade-off that separates species with fast growth from species with high survival and a stature–recruitment trade-off that separates species that achieve large stature from species with high recruitment. As these trade-offs have typically been explored for trees ≥ 1 cm dbh, it is unclear how species' growth and survival during earliest seedling stages are related to the trade-offs for trees ≥ 1 cm dbh. Here, we used principal components and correlation analyses to (1) determine the main demographic trade-offs among seed-to-seedling transition rates and growth and survival rates from the seedling to overstory size classes of 1188 tree species from large-scale forest dynamics plots in Panama, Puerto Rico, Ecuador, Taiwan, and Malaysia and (2) quantify the predictive power of maximum dbh, wood density, seed mass, and specific leaf area for species' position along these demographic trade-off gradients. In four out of five forests, the growth–survival trade-off was the most important demographic trade-off and encompassed growth and survival of both seedlings and trees ≥ 1 cm dbh. The second most important trade-off separated species with relatively fast growth and high survival at the seedling stage from species with relatively fast growth and high survival ≥ 1 cm dbh. The relationship between seed-to-seedling transition rates and these two trade-off axes differed between sites. All four traits were significant predictors for species' position along the two trade-off gradients, albeit with varying importance. We concluded that, after accounting for the species' position along the growth–survival trade-off, tree species tend to trade off growth and survival at the seedling with later life stages. This ontogenetic trade-off offers a mechanistic explanation for the stature–recruitment trade-off that constitutes an additional ontogenetic dimension of life-history variation in species-rich ecosystems.

KEYWORDS

Barro Colorado Island, ForestGeo forest dynamics plot, Fushan, life-history strategy, Luquillo, Pasoh, seed size, species coexistence, Yasuní

INTRODUCTION

Following the general principles of life-history theory, all organisms must partition limited resources among demographic processes related to growth, survival, and reproduction, which leads to trade-offs between demographic rates that facilitate the local coexistence of species with different life-history strategies (Lande, 1982; Stearns, 1992). On the global scale, the diversity of plant life-history strategies tend to be aligned along two main gradients: the fast–slow continuum (Reich, 2014) and a gradient in plant reproductive strategies (Salguero-Gómez et al., 2016).

On the local scale, the study of simultaneous trade-offs among coexisting and competing life-history strategies requires large diversity of co-occurring species, such as can be found in old-growth grasslands (Lind et al., 2013; Nerlekar & Veldman, 2020) and tropical forest ecosystems which can harbor up to hundreds of coexisting tree species and life-history strategies (Condit et al., 2006; Ohse et al., 2023; Worthy & Swenson, 2019).

Previous studies have shown that the local diversity of life-history strategies in tropical forests is often structured along two orthogonal demographic trade-offs: the well-known growth–survival trade-off which separates

species with fast growth rates from species with high survival rates (the fast–slow continuum) and the less well-studied stature–recruitment trade-off which separates species with tall adult stature from species with high recruitment success (Kambach et al., 2022; Kohyama, 1993; Rüger et al., 2018). However, most of these previous analyses have only included growth and survival of trees ≥ 1 cm diameter at 1.3 m above ground (dbh, Russo et al., 2021; Wright et al., 2010), thus neglecting how the growth and survival rates of seedlings relate to growth and survival at larger tree size classes.

At Barro Colorado Island (BCI) in Panama, seedling growth and survival were related to both demographic trade-offs (Rüger et al., 2018); that is, most species that grow fast as seedlings also tend to grow fast as adults while species that survive well as seedlings also achieve high survival at larger size classes (Gilbert et al., 2006; Kunstler et al., 2009). Yet, after accounting for this general trade-off between growth and survival rates at BCI, the remaining variation in life-history strategies was structured along an orthogonal gradient that separated species with high recruitment rates and above-average growth and survival rates at the seedling stage from species with tall adult stature and above-average growth and survival rates at larger size classes (i.e., ≥ 1 cm dbh, King et al., 2006; Rüger et al., 2018). However, it is not known whether the observed association between the stature–recruitment trade-off and a trade-off in growth and survival rates between seedlings and larger size classes observed at BCI holds at other forest sites.

Another limitation in studies of life-history strategies in forest ecosystems relates to the quantification of reproduction, which is often represented by recruitment rates over a given stem size threshold (e.g., ≥ 1 cm dbh). To derive a comparable measure of allocation to reproduction among species, the number of recruits needs to be scaled by either the number of seeds produced, the resources invested into seedling survival, or the number of reproductive trees (Bogdziewicz et al., 2023). Since information on the reproductive status of trees or production of seeds is not often available and allocation to seedling survival cannot be measured at all, recruitment rates can sometimes only be scaled by the number or basal area of conspecific “presumably reproductive” trees (e.g., Kambach et al., 2022; Rüger et al., 2018). This scaling approach hinges on the uncertainty about the size at which species become reproductive and bears the potential to neglect differences between monoecious and dioecious species. To avoid this problem, we collated data on seed production from seed traps and data on seedling establishment from seedling censuses to calculate seed-to-seedling transition rates, a measure of reproductive success that does not require any scaling. This allowed us to quantify

how seed-to-seedling transition rates were related to different axes of life-history differentiation (Muscarella et al., 2013) and whether seed-to-seedling transition rates showed a negative relationship with species maximum stature as predicted by Kohyama (1993).

In trees (as in plant species in general), species’ life-history strategies and thus their position along the demographic trade-off gradients, are manifested through differences in key functional traits such as plant stature, stem or wood density, seed mass, and leaf specific area (Adler et al., 2014; Díaz et al., 2016). At BCI (Rüger et al., 2018), species’ positions along the growth–survival trade-off gradient were predicted by differences in wood density and seed mass (with lower wood density and lower seed mass being indicative of faster life-history strategies), while species’ positions along the stature–recruitment trade-off gradient were predicted by differences in maximum height and seed mass (with larger maximum height and seed mass being indicative of lower recruitment rates). While these trait-demography relationships also tend to hold at the seedling stage (Metz, Wright, et al., 2023) their relative importance can shift with ontogenetic stage (Visser et al., 2016) and we lack a comprehensive understanding of the predictive value of plant traits for life-history variation across all size classes. An understanding of these trade-offs is urgently needed to accurately assign tree species into groups of similar form and life-history strategy (Rüger et al., 2018), which can then be used to predict forest dynamics (Rüger et al., 2020) and species’ vulnerability to climate- and land-use change (Ohse et al., 2023).

In this study, we compiled seed, seedling, and tree census data from five tropical forests in Asia and the Americas to calculate species-level seed-to-seedling transition rates as well as growth and survival rates in two seedling size classes (from seed germination < 20 cm and 20–50 cm height) and two tree size classes that reflect differences in light availability (overstory, understory). Using principal components and post hoc correlation analyses, we identified the trade-offs that structure the local diversity in life-history strategies and their relationships with four key functional traits (maximum dbh, wood density, seed mass, and specific leaf area). We investigated (1) how seedling growth, survival and seed-to-seedling transition rates relate to the demographic rates of larger trees and (2) how the four functional traits relate to the main demographic trade-offs across the full-size spectrum of trees. Specifically, we asked whether seedling growth and survival rates align with the growth–survival trade-off for larger trees and to what degree is the stature–recruitment trade-off associated with a growth or survival trade-off between ontogenetic

stages (i.e., between seedlings and larger tree size classes). The results provided us with a better understanding of the demographic trade-offs that structure local plant diversity across the full-size spectrum and improved our understanding of the mechanisms that facilitate species coexistence in diverse plant communities.

METHODS

Study sites

We assembled data on freestanding woody plant species from five tropical forest dynamics plots of 16–50 ha from the ForestGeo network (Anderson-Teixeira et al., 2015; Davies et al., 2021). The five sites are BCI (Panama), Fushan (Taiwan), Luquillo (Puerto Rico), Pasoh (Malaysia), and Yasuní (Ecuador) (Table 1). The forest at Luquillo experienced major cyclonic storms in 1989 and 1998. The forest at Fushan is exposed to major storms annually during the typhoon season from July to September each year and experienced major cyclonic storms in 2004, 2008, and 2013 (Hogan et al., 2018). At Pasoh, the forest understorey was heavily disturbed by a 100-fold increases in native wild boar populations, which fed on nearby oil-palm plantations and used the understorey plants to construct their birthing nests (Luskin et al., 2017). The forests at BCI and Yasuní had no major disturbances during the study period. In each forest, individuals <1 cm dbh (dbh 1.3 m from the ground, hereafter seedlings) were mapped and measured in subplots of the ForestGeo plots, and all individuals ≥1 cm dbh (hereafter trees) were mapped and measured following Condit

(1998). All five ForestGeo plots have been re-censused approximately every five years, with different time periods used for this study, as a result of data availability (BCI: 1982–2015, Yasuní: 1996–2008, Pasoh: 1987–2010, Luquillo: 1992–2016, and Fushan: 2004–2013).

For all five forests, we combined seed trap data with recruitment data from annual seedling censuses to calculate seed-to-seedling transition rates, and we assembled data on species' maximum dbh, wood density, and specific leaf area. In addition, we assembled data on species seed mass (these data were not available for Pasoh). In the following sections, we have described the measurements and calculations of all species-level demographic rates and traits.

Data on growth and survival of trees ≥1 cm dbh

In every census, each individual with at least one living stem ≥1 cm dbh was identified to species or morphospecies, determined as “alive” and the stem with the largest dbh was measured with an accuracy of 1 mm following the protocol presented in Condit (1998). For each individual tree where the main stem survived a census interval, we calculated growth as the absolute annual increment in dbh $\frac{dbh_{t2} - dbh_{t1}}{t}$ with t = time in years between consecutive censuses.

To account for the dependence of growth and survival on tree size and light availability, we assigned each tree to the overstory or the understory layer, based on the perfect plasticity approach as described in Kambach et al. (2022).

TABLE 1 Characteristics of the five long-term forest dynamics plots.

ForestGeo plot (country)	Latitude; longitude	MAT (°C)	Precipitation (mm)	Plot size (ha)	Census dates	Forest type (disturbance type)	Species no.	Allometric coefficients [†]	
								<i>a</i>	<i>b</i>
Barro Colorado Island, BCI (Panama)	9°9'15.48"; −79°50'45.96"	27.1	2551	50	1982, 1985, 1990, 1995, 2000, 2005, 2010, 2015	Broadleaf evergreen (seasonal drought)	299	0.03615016	1.281928
Fushan (Taiwan)	24°45'41.04"; 121°33'18"	18.2	4271	25	2004, 2008, 2013	Broadleaf evergreen (cyclonic storms)	101	0.01652303	1.3
Luquillo (Puerto Rico)	18°19'34.32"; −65°48'57.6"	22.8	3548	16	1992, 1996, 2001, 2006, 2011, 2016	Broadleaf evergreen (cyclonic storms)	138	0.02477302 [‡]	1.320 [‡]
Pasoh (Malaysia)	2°58'55.2"; 102°18'46.8"	27.9	1788	50	1987, 1990, 1995, 2000, 2005, 2010	Mixed evergreen dipterocarp (everwet)	814	0.1539492	1.204044
Yasuní (Ecuador)	0°41'9.244"; −76°23'49.2"	28.3	3081	50	1996, 2003, 2008	Broadleaf evergreen (everwet)	1114	0.03615016	1.281928

Abbreviation: MAT, mean annual temperature.

[†]To calculate crown area (in square meters) = $a \times dbh$ (in millimeters)^b.

[‡]Separate coefficients for Arecaceae with $a = 0.002200656$, $b = 1.647$.

The crown area of each individual tree was approximated with the allometric equation crown area (in square meters) = $a \times \text{dbh}$ (in millimeters)^{*b*}. For each species, the allometric coefficients *a* and *b* were either calculated from the Tallo global tree allometry and crown architecture database (for species with ≥ 10 observations, Jucker et al., 2022) or assigned from site-specific allometries (listed in Table 1). For BCI and Yasuní, the site-specific coefficients were extracted from Bohlman and Pacala (2012). For Fushan, the coefficients were calculated from 1228 individuals from 65 tree species (P.J. Luo & C.-H. Chang-Yang, unpublished data). For Luquillo, separate coefficients for dicots and monocots were derived from the equations presented in Zambrano et al. (2019). For Pasoh, coefficients were calculated from the six largest individuals from the 100 most abundant species in the 2005 census (S. J. Wright, unpublished data). Each ForestGeo plot was divided into subplots of 31.5×31.5 m (following Bohlman & Pacala, 2012), in which we ranked all individuals from the largest to the smallest diameter. We successively assigned the largest individuals to the overstory canopy layer, until their cumulative crown area exceeded twice the subplot area. The remaining smaller individuals were then assigned to the understory canopy layer. This means that trees assigned to the overstory are expected to be shaded by, at a maximum, one other tree crown, whereas trees assigned to the understory are shaded by more than two other tree crowns. Palms (Arecaceae), hemi-epiphytic, and unidentified species were retained for assignment to the overstory or understory layer but were omitted from all other analyses (this excluded 15 species at BCI, 3 at Fushan, 3 at Luquillo, 46 at Pasoh and 26 at Yasuní).

Data on growth and survival of seedlings ≤ 50 cm height

At all five sites, growth, survival, and recruitment rates of seedling individuals < 1 cm dbh were measured according to ForestGeo protocols (described in Davies et al., 2021); that is, each seedling was individually tagged, identified to species or morphospecies, deemed dead, missing, or alive, and (in the latter case) its height was measured with an accuracy of up to 1 mm. As our study focused on growth and survival rates during earliest life stages, we only calculated growth and survival rates for two seedling classes: individuals < 20 cm height and individuals of 20–50 cm height. Size classes were assigned based on seedling height at the first of two consecutive censuses and independent from seedling height at the second census.

For BCI, we combined measurements from two different datasets. Seedlings < 20 cm height were censused

annually from 1994 to 2019 (omitting measurements that were more than 1.5 years apart) in 600 seedling plots of 1 m^2 (Wright et al., 2005, 2015) and seedlings of ≥ 20 cm height were censused annually from 2001 to 2004, 2008 to 2009, and 2011 to 2013 in $\sim 20,000$ evenly stratified 1-m^2 seedling plots (Comita et al., 2007, 2023). At Fushan, seedlings were censused quarterly from 2003 to 2010 in 264 seedling plots, and biannually from 2011 to 2019 in 318 seedling plots of 1 m^2 , each one located at a distance of 2 m from 106 focal seed traps (Chang-Yang et al., 2013). At Luquillo, seedlings were censused annually from 2007 to 2016 in 360 seedling plots of 1 m^2 , each one located at a distance of 2 m around 120 focal seed traps (Muscarella et al., 2013; Zimmerman, 2018). At Pasoh, seedlings were censused annually from 2001 to 2008 in up to 1010 seedling plots of 1 m^2 (some extra seedling plots were added during this time range), each one located at a distance of 2 m around 336 focal seed traps (Chen, 2007). At Yasuní, seedlings were censused annually from 2002 to 2019 in 600 seedling plots of 1 m^2 , each one located at a distance of 2 m around 200 focal seed traps (Metz et al., 2010).

For each individual seedling that survived a census interval, we calculated growth as the absolute annual increment in height $\frac{\text{height}_{t_2} - \text{height}_{t_1}}{t}$ with *t* = time in years between consecutive censuses.

Data on seed production

Seed fall was recorded in seed traps according to ForestGeo protocols that are described in Davies et al. (2021). At BCI, seeds were collected weekly from 1993 to 2012 in 200 traps of 0.5 m^2 (Visser et al., 2016; Wright et al., 2005). At Fushan, seeds were collected monthly from 2002 to 2007 in 87 traps and from 2007 to 2019 in 106 traps of 0.5 m^2 (Chang-Yang et al., 2024). At Luquillo, we used seed trap data that were collected biweekly from 1993 to 2006 in traps of 0.16 m^2 and biweekly from 2007 to 2021 in 121 traps of 0.5 m^2 (shifted to accommodate ForestGeo protocols). At Pasoh, seed were collected weekly from 2001 to 2020 in traps of 0.5 m^2 . We only included data from 2002–2010, 2012–2014, and 2018 as the other years had > 2 weeks of missing seed rain observations. At Yasuní, seeds were collected twice each month from 2000 to 2018 in 200 traps of 0.5 m^2 (Metz et al., 2010).

Data on species traits

Species-level trait data were assembled from primary and published sources as follows. For BCI, wood density, specific leaf area, and maximum height measurements

are described in Wright et al. (2010) and Rüger et al. (2018). Mean dry seed mass was estimated by averaging measurements from up to five seeds from up to five fruits from up to five individuals in the ForestGeo plot or adjacent areas (Metz et al., 2023; Wright et al., 2010), supplemented with data from Daws et al. (2005). For Fushan, wood density was obtained from plot-level measurements and from the literature (Chave et al., 2009; Lasky et al., 2013), mean dry seed mass was measured from the seed rain data or obtained from the literature, and specific leaf area was measured on 6–12 individuals of each species and maximum height was measured according to established protocols (Cornelissen et al., 2003). For Luquillo, wood density, mean dry seed mass, specific leaf area, and maximum height measurements are described in Swenson et al. (2012) and Umaña et al. (2016). At Pasoh, wood density for 388 species was extracted from the global wood density database (Chave et al., 2009; Zanne et al., 2009) and specific leaf area and mean maximum height were calculated for the 100 most abundant species (in 2005), as measured from the six largest individuals (based on dbh). At Yasuní, mean dry seed mass was estimated from a pooled sample of seeds from fruiting individuals or seed traps (Fortunel et al., 2016; Metz et al., 2023). Wood density was measured for 2052 trees in 2010 and covers three measurements for 558 species, two measurements for 34 species and single measurements for 16 species. Specific leaf area measurements are described in Kraft et al. (2008).

Maximum dbh was calculated as the mean dbh of the six largest trees of each species at the respective ForestGeo plot. To test if maximum dbh is an appropriate proxy for stature, we tested the relationship between maximum dbh and maximum height in BCI, Fushan, Luquillo, and Pasoh using the Pearson correlation coefficient.

Calculation of species-specific demographic rates

For each of the five forests, we estimated species-specific mean annual growth and survival rates in four size and light availability classes (seedlings <20 cm height, seedlings 20–50 cm height, trees ≥1 cm dbh in the understory and in the overstory canopy) as well as species-specific mean seed-to-seedling transition rates. We only calculated growth and survival rates for species and size classes when there were ≥10 observations.

To speed up computations and to linearize the relationships between variables for principal components analysis, we transformed the observed annual growth rates to an approximate normal distribution using the Modulus

transformation with $\lambda = 0.4$ (Equation 1) as described in John and Draper (1980) and Condit et al. (2017).

$$\text{growth}_{\text{trans}} = \begin{cases} \text{growth}^\lambda & g \geq 0 \\ -\{(-\text{growth})^\lambda\} & g < 0 \end{cases} \quad (1)$$

To calculate species-specific growth rates within each size class, we randomly selected (without replacement) up to 200 growth observations across all census intervals (to speed up computation time) per species and size class. We used hierarchical Bayesian models to calculate the mean transformed growth rate g of species j in size class k with the likelihood of the observed transformed growth rate $\text{growth}_{i,j}$ of individual i of species j in size class k as follows (Equation 2):

$$\begin{aligned} \text{growth}_{i,j,k} &= \text{Normal}(g_{j,k}, \text{sigma}_{j,k}) \\ g_{j,k} &= \text{Normal}(g_k, \sigma_{g,k} = 1) \\ \text{sigma}_{j,k} &= \text{Lognormal}(\text{sigma}_k, \sigma_{\text{sigma},k} = 1) \end{aligned} \quad (2)$$

with g_k and sigma_k being the size class-specific average transformed growth rate (across all species) and respective SD (both with uniform flat priors).

To calculate species-specific survival rates within each size class, we randomly selected (without replacement) up to 1000 survival observations across all census intervals (to speed up computations) per species and size class. We used hierarchical Bayesian models with flat priors to calculate the mean annual survival rate s of species j with the likelihood of the observed individual i of species j in size class k remaining alive $i_{j,k}$ between consecutive censuses as follows (Equation 3):

$$\begin{aligned} p(\text{alive}_{i,j,k} | s_{j,k}) &= \text{Bernoulli}(\text{alive}_{i,j,k} | s_{j,k}^{t_i}), \\ s_{j,k} &= \text{Beta}(\alpha_k, \beta_k) \end{aligned} \quad (3)$$

with α_k and β_k being the size class-average parameters of the beta distribution. We transformed survival rates to the logarithm of the expected lifespan via $\log\left(\frac{1}{1 - \text{survival probability}}\right)$ (based on constant survival probabilities, c.f. Condit, 2022).

To calculate comparable seed-to-seedling transition rates, we first calculated, for each forest and census interval, the annual number of newly emerging seedlings per hectare (scaled from the summed area of seedling plots) and the annual number of seeds captured per hectare (scaled from the summed area of seed traps). Seed-to-seedling transition rates were then calculated by dividing

the number of newly emerged seedlings per hectare per year by the number of seeds captured per hectare per year and averaged across all census intervals. Mean seed-to-seedling transition rates >1 were replaced with a value of one. This applied to two species at Luquillo (*Guarea glabra* Vahl, *Syzygium jambos* (L.) Alston), one species at Pasoh (*Xerospermum noronhianum* Blume), and three species at Yasuní (*Clarisia biflora* Ruiz & Pav., *Compsonura capitellata* (Poepp. ex A. DC.) Warb., and *Crepidosperrum rhoifolium* Triana & Planch). We then log-transformed seed-to-seedling transition rates.

Statistical analyses

We normalized each demographic rate to unit variance (via subtraction of the mean and division by the SD, separately per size class and ForestGeo plot) and used principal components analyses to analyze their joint relationships (both within and across forest sites).

Most species did not have data for all four size classes (Appendix S1: Figure S1), which is a prerequisite for traditional principal components analysis. To be able to include as many species as possible, we retained all species with growth and survival estimates in at least two size classes and used an iterative algorithm to impute missing growth and survival estimates with values that guaranteed a minimized effect on the first four principal components of the available observations (Josse & Husson, 2016). This procedure provides principal components that are robust to the imputation of missing values (Appendix S1: Figures S2–S6).

To test the significance of the resulting principal components, we compared the observed eigenvalues to a distribution of eigenvalues obtained from 1000 random permutations of the demographic rates across species (i.e., against null datasets with no correlation structure, Camargo, 2022). Factors with a loading value of ≥ 0.3 were determined to be significantly related to the respective principal component. Principal components with significant loadings of growth versus survival rates in opposite directions were considered to capture the growth–survival trade-off. Principal components with significant loadings of seedling versus tree growth rates or seedling versus tree survival rates in opposite directions were considered to capture the stature–recruitment trade-off in demographic rates.

For each ForestGeo plot, we quantified the relationships among species functional traits (maximum dbh, wood density, specific leaf area, and seed mass) and the main trade-offs between growth, survival, and seed-to-seedling transition rates (represented by the first

two principal components). As trait information were not available for all species (Appendix S1: Figure S1), we determined the significance of these relationships with separate post hoc correlation tests for each trait and applied a Bonferroni-adjusted significance threshold of $p < 0.0125$.

For each pair of ForestGeo plots, we tested for the similarity in principal components by means of a Procrustes analysis (Legendre & Legendre, 2012), a statistical shape analysis that determines optimal scale, shift, and rotation parameters to compare the distribution of two sets of points in n -dimensional space. Here, we used Procrustes analyses to test the multivariate correlation (at $p < 0.05$) between the factor loadings of species' demographic rates along the first two principal components between two forest sites.

To quantify global relationships and life-history variation among the five investigated forests, we conducted a global principal components analysis in which the contribution of each species was weighted according to the inverse of the local species number (to guarantee an equal contribution of each ForestGeo plot). The same imputation method (for dealing with missing data) and significance tests were applied as for the single site analyses (outlined above). Using species' scores along the first and second principal component, we determined the core position and size of the site-specific local life-history spaces with minimum convex polygons that covered 10% and 90% of co-occurring species, respectively.

All analyses were conducted in R, version 4.2.2 (R Core Team, 2021), using the following R-packages: rstan and coda to run Bayesian hierarchical models (Plummer et al., 2006; Stan Development Team, 2018), missMDA and PCAtest for data imputation and principal components analysis (Camargo, 2022; Josse & Husson, 2016), ggplot2 for graphical representations (Wickham, 2009), and vegan for post hoc correlation and centroid calculations (Oksanen et al., 2020).

RESULTS

Growth rates were positively correlated for seedlings and individuals ≥ 1 cm dbh and survival rates were positively correlated for seedlings and individuals ≥ 1 cm dbh in three of the five forests (BCI, Pasoh, and Yasuní, Figure 1; pair-wise correlations are shown in Appendix S1: Figures S7–S11). At Luquillo, seedling growth was likewise positively related to growth of trees ≥ 1 cm dbh, but survival of seedlings < 20 -cm and 20–50-cm tall was negatively related to survival in the highest canopy layers. At Fushan, however, seedling growth was negatively related to growth of trees ≥ 1 cm dbh and

seedling survival was negatively related to survival in the highest canopy layers.

At each of the five forests, the first two principal components captured a significant proportion of variation in demographic rates and accounted for 56.3%–80.4% of the joint variation in growth, survival, and seed-to-seedling transition rates (Figure 1).

Except for Fushan, the first principal component was associated with a general trade-off between growth and survival rates (growth–survival trade-off, Figure 1) and the second principal component was associated with a trade-off between either growth rates, survival rates, or both for the seedling versus tree size classes ≥ 1 cm dbh (Figure 1). At Fushan, the relative importance of the two trade-off gradients was switched so that the trade-off between growth and survival rates at the seedling versus larger tree size classes ≥ 1 cm dbh was associated with the first principal component and the growth–survival trade-off was associated with the second principal component.

Across the five forests, seed-to-seedling transition rates were consistently positively related to the survival rates of seedlings height < 20 cm but showed otherwise varying relationships with species' demographic rates (Appendix S1: Figures S2–S6). Along the principal components of demographic variation, seed-to-seedling transition rates were significantly associated with the growth–survival trade-off at BCI, Fushan, Luquillo, and Pasoh (positively correlated with species' survival rates) and significantly associated with the stature–recruitment trade-off at BCI, Fushan, Pasoh, and Yasuni (positively correlated with seedling growth and survival rates).

Regarding the similarity in principal components among the five forests, pair-wise Procrustes analyses showed that the positions of species' demographic rates along the first two principal components were similar among all five forests (with r ranging from 0.77 to 0.98 and p ranging from 0.008 to < 0.001).

The four traits (maximum dbh, wood density, seed mass, and specific leaf area) emerged as significant predictors of species positions along the first two principal components of demographic variation (Figure 2) in all five forests, except for Pasoh where specific leaf area was unrelated to the first two principal components. The relative predictive power of the four traits (inferred from the % of explained variation) differed among the five forests. Maximum dbh (a proxy for maximum height, Appendix S1: Figure S12) accounted for 14%–63% of demographic variation and was positively related to the growth and survival rates of trees ≥ 1 cm and negatively related to the survival rates of seedlings (Appendix S1: Figures S7–S11). Accordingly, maximum dbh was associated with the trade-off between growth and/or survival at the seedling

versus larger size classes of trees ≥ 1 cm dbh, indicating a trade-off between maximum dbh and seedling demographic rates (i.e., the stature–recruitment trade-off). Wood density accounted for 11%–30% of demographic variation and was most strongly, and positively, related to survival rates ≥ 1 cm dbh in all five forests. Specific leaf area accounted for 3%–27% of demographic variation and was positively related to seedling growth but negatively related to maximum dbh across all five forests. Seed mass accounted for 10%–46% of demographic variation and positively related to seed-to-seedling transition rates and survival of seedlings < 20 -cm tall in all four forests with available data (Appendix S1: Figures S7–S9 and S11).

When all species- and site-specific demographic rates were analyzed in a global principal components analysis (with equal weights among forests sites), the resulting first and second principal components accounted for 70.7% of the global demographic variation and were associated with the growth–survival and stature–recruitment trade-offs, respectively (Figure 3). Seed-to-seedling transition rates were closely aligned with higher survival along the growth–survival trade-off. The site-specific 10% core life-history spaces mostly aligned along the growth–survival trade-off, and the 10% core spaces of Pasoh and Fushan were completely separated from the core spaces of BCI, Luquillo and Yasuni. Life-history spaces that included 90% of all species-specific life-history characteristics, however, largely overlapped among the five forests.

DISCUSSION

In this study, we investigated the main demographic trade-offs that structured the spectra of life-history strategies from seedlings to the largest size classes of 1188 tree species in five tropical forests. In four out of the five forests, the growth–survival trade-off for seedlings and trees was the most important demographic trade-off (accounting for about 50% of demographic variation, Kambach et al., 2022; Rüger et al., 2018; Russo et al., 2021). After accounting for the growth–survival trade-off, that is, when comparing only species that had a similar position along the fast–slow continuum, we observed that these species trade-off investment between growth and/or survival at early life stages (seedlings) versus investment at later life stages (trees). The emergent ontogenetic trade-off between different life stages provides a mechanistic explanation for the previously reported stature–recruitment trade-off (Kambach et al., 2022; Rüger et al., 2018), that is, the second orthogonal dimension of life-history differentiation that separates species with relatively fast growth and high survival at the seedling stage from species that are able to achieve a large




FIGURE 1 Principal components analyses on species-specific growth, survival, and seed-to-seedling transition rates of tree and shrub species in five tropical forests. Depicted are the joint (arrows) and separate factor loadings (bars) along the first and second principal component. Asterisks indicate significant principal component loadings following permutation tests at $p < 0.05$ (Josse & Husson, 2016).




FIGURE 2 Post hoc correlation tests of four functional traits (black) with the factor loadings obtained from separate principal components analyses of species-specific growth, survival, and seed-to-seedling transition rates (colored) of woody tree and shrub species in five tropical forests (c.f. Figure 1). Black arrows show linear 2D trend surfaces, that is, indicate the direction of the steepest increase in trait values (scaled to unit length) along the first two principal components. Dotted arrows indicate non-significant relationships between principal components and traits. Text labels show the proportion of explained variation, the number of species, and the Bonferroni-corrected significance of the post hoc correlation at $p < 0.015$.




FIGURE 3 Global principal components analysis and life-history spaces across the growth, survival, and seed-to-seedling transition rates of 1188 tree and shrub species in five tropical forests. Depicted are the joint factor loadings (arrows) and species positions (dots) along the first and second principal component together with the minimum convex hulls that cover the 90% area (dashed lines) and 10% core area (shaded areas) of site-specific life-history spaces. Asterisks on axes labels indicate significant principal component loadings following permutation tests at $p < 0.05$ (Josse & Husson, 2016).

maximum stature (Kambach et al., 2022). Relationships of seed-to-seedling transition rates with the observed demographic trade-offs, however, were inconclusive.

Beyond our model system of tropical forests, the growth–survival trade-off (i.e., the fast–slow continuum of life-history strategies) has previously been evaluated as a general feature that structures the diversity of plants (Roberto Salguero-Gómez et al., 2016), fish (Beukhof et al., 2019), mammals (Oli, 2004), and arthropods (Bakewell et al., 2020; Blackburn, 1991). In grasslands, a similar trade-off can be observed between fast growth versus increased defense against consumers and pathogens (He et al., 2022; Lind et al., 2013). Trade-offs among growth or survival rates between ontogenetic stages or size classes, however, have less often been investigated. Although pleiotropy theory predicts that finite populations

should be structured by a selective pressure that enforces a trade-off between juvenile and adult survival (Giaimo, 2014), such a trade-off between ontogenetic stages might be difficult to quantify as it could be masked by the generally stronger signal of the slow–fast continuum. In addition, the relative ecological strategies of plants species are hypothesized to shift with ontogeny-dependent filters (Dayrell et al., 2018; Wijenayake et al., 2023).

In the cyclonic storm-disturbed forests at Luquillo and Fushan, the growth–survival trade-off involved only survival at the seedling stage (Luquillo) or understory size class (Fushan). In contrast to previous studies that focussed on trees ≥ 1 cm dbh or contrasted growth in high-resource conditions versus survival in low-resource conditions (Kambach et al., 2022; Russo et al., 2021;

Umaña et al., 2023), the inclusion of seedling census data showed that even the two cyclonic storm-impacted forests were structured by a growth–survival trade-off (c.f., Metz et al., 2023) and a stature–recruitment trade-off (described as a fecundity–stature axis by Umaña et al., 2023). The life-history strategies in these two cyclone-disturbed forests exhibited a negative relationship between survival rates in the seedling stage versus survival rates in the highest canopy layers (described as a survival axis by Umaña et al., 2023). These results highlight that cyclonic storms can condition trade-offs such as between competitive and colonization ability that are usually not observed in less disturbed tropical forest communities (Kambach et al., 2022; Russo et al., 2021; Umaña et al., 2023; Uriarte et al., 2012).

Although seed-to-seedling transition rates showed a consistently positive relationship with seedling survival <20 cm height, the relationships with other demographic rates and trade-off gradients were inconsistent among the five forests studied. In Pasoh, which is the forest most strongly affected by unnaturally high wild boar populations that caused much disturbance to seedlings in the understorey (Ickes & Thomas, 2003; Luskin et al., 2017), seed-to-seedling transition rates were largely decoupled from demographic rates (but not species traits). In the other four forests, we found a negative relationship between maximum stature and seed-to-seedling transition rates, which is in line with a mechanism proposed by Kohyama (1993) that enables tree species coexistence in forests with pronounced gap dynamics. We found that species with the fastest growth rates had the lowest seed-to-seedling transition rates (except at Fushan), which is consistent with higher seedling mortality associated with smaller seed sizes observed at BCI, Luquillo, and Yasuní (Muscarella et al., 2013; Wright, Calderón, et al., 2024).

In accordance with previous studies, the four plant traits (maximum dbh, wood density, seed mass, and specific leaf area) emerged as significant predictors for the life-history strategies of tree species (Metz et al., 2023; Rüger et al., 2018; Wright et al., 2010) and explained a comparatively large proportion of demographic variability (c.f. Paine et al., 2015). The specific relationships between traits and growth and survival rates depended strongly on the ontogenetic stage (especially for specific leaf area, seed mass, and maximum height and less so for wood density, Gibert et al., 2016; Lasky et al., 2015). Since leaf and wood traits used for this analysis were measured from individuals ≥ 1 cm dbh, this could have reduced the strength of their relationships with demographic rates at seedling size classes.

Although our results are based on one of the most comprehensive datasets available on hyperdiverse tree

communities, the observed trade-offs are derived for the rather common species and might have missed fast-growing species that died or outgrew the 50-cm seedling height threshold between censuses. One potentially important dimension of life-history differentiation that we did not explicitly consider includes any potential trade-offs with fecundity (i.e., biomass production of flowers, fruits, and seeds, Muller-Landau, 2010; Villellas & García, 2013). Quantifying any trade-off with the different components of reproduction is still difficult because reproduction effort must be standardized by the local abundance, basal area, or crown area of reproductive individuals (c.f. Kambach et al., 2022) which, to our knowledge, has only been done at BCI (Visser et al., 2016). Here, the Masting Inference and Forecasting network offers a great opportunity to study the global trade-offs between seed size, seed mass, standardized fecundity, and tree senescence (MASTIF, Qiu et al., 2021, 2022). Finally, our sample size of five forests was too low to investigate how the growth–survival or the stature–recruitment trade-off might be affected by soil nutrients (Russo et al., 2008), precipitation (Browne et al., 2023), elevation (de Kamimura et al., 2023), biogeographic region (Kambach et al., 2022), or disturbance regimes (Russo et al., 2021). To provide more generalizable patterns across global gradients in climate and biogeography, we encourage the dedicated collection of seedling demographic data with standard protocols in disturbed and undisturbed forests (Davies et al., 2021).

In conclusion, our study adds to the growing body of cross-site analyses on the trade-offs that structure the demographic diversity forest ecosystems (Condit et al., 2006; Kambach et al., 2022; Needham et al., 2022; Russo et al., 2021). Beside the well-documented trade-off between fast growth and high survival rates, we identified an ontogenetic dimension of life-history variation that may contribute to coexistence between species that prioritize growth and/or survival at the seedling/juvenile stage and species that prioritize growth and survival at larger size classes. Based on observations from five tropical forests, we hypothesize that these two trade-offs could pose two general features that structure diverse plant communities.

AUTHOR CONTRIBUTIONS

Stephan Kambach contributed to conceptualization, data curation, formal analysis, investigation, software, visualization, and writing—original draft preparation. Helge Bruelheide contributed to funding acquisition, conceptualization, data analysis, writing—review and editing. Liza S. Comita contributed to funding acquisition, conceptualization, investigation, writing—review and editing. Richard Condit contributed to conceptualization, investigation,

writing—review and editing. Salomón Aguilar, Chia-Hao Chang-Yang, Pei-Jen Luo, Rolando Pérez, Simon A. Queenborough, Nathan G. Swenson, and Tze Leong Yao contributed to investigation, writing—review and editing. Yu-Yun Chen, Nancy C. Garwood, Stephen P. Hubbell, Margaret R. Metz, Musalmah Bt. Nasardin, I-Fang Sun, María Uriarte, Renato Valencia, S. Joseph Wright, and Jess K. Zimmerman contributed to funding acquisition, investigation, writing—review and editing. Jill Thompson contributed to funding acquisition, investigation, data curation, writing—review and editing. Nadja Rüger contributed to conceptualization, data curation, funding acquisition, methodology, supervision, writing—review and editing.

ACKNOWLEDGMENTS

We are indebted to all the people involved in the establishment and ongoing measurement at the ForestGEO research sites. We would like to acknowledge the general support from the Forest Global Earth Observatory (ForestGEO), the Center for Tropical Forest Science and the Smithsonian Tropical Research Institute in Panama, the Japan Society for the Promotion of Science (JSPS), the National Science Foundation (NSF), the National Geographic Society (NGS), and the Conservation, Food and Health Foundation (CFHF). The work of Stephan Kambach was funded by the iDiv-Flexpool—the internal funding mechanism of the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, a research center of the German Research Foundation (DFG—FZT 118) and by the 2019–2020 BiodivERsA joint call for research proposals, under the BiodivClim ERA-Net COFUND program with the funding organizations Swiss National Science Foundation SNF (FeedBaCks 193907), Agence Nationale de la Recherche (ANR-20-EBI5-0001-05), the German Research Foundation (DFG BR 1698/21-1, DFG HI 1538/16-1), and the Technology Agency of the Czech Republic (SS70010002). Stephan Kambach acknowledges the support of the Smithsonian Tropical Research Institute to attend the ForestGEO analytical workshop in 2018 and 2019. The BCI forest dynamics research project (Condit, 1998; Condit et al., 2019; Hubbell et al., 1999) was founded by S.P. Hubbell and R.B. Foster and is now managed by R. Condit, S. Lao, and R. Perez. Numerous organizations have provided funding, principally the U.S. National Science Foundation (NSF). Seedling data collection in the BCI plot were funded by multiple NSF LTREB grants (DEB 0075102, DEB 0823728, DEB 0640386, DEB 1242622, and DEB 1464389). The Fushan forest dynamics plot (Su, 2007) is supported by the Taiwan Forestry Bureau, the Taiwan Forestry Research Institute and the Ministry of Science and Technology of

Taiwan. We also thank the staff at Fushan Research Center for providing logistic support. The Luquillo forest dynamics plot was supported by grants BSR-8811902, DEB 9411973, DEB 0080538, DEB 0218039, DEB 0620910, DEB 0963447, and DEB-129764 from the NSF to the Department of Environmental Science, University of Puerto Rico, and to the International Institute of Tropical Forestry, USDA Forest Service, as part of the Luquillo Long-Term Ecological Research Program. The U.S. Forest Service (Dept. of Agriculture), and the University of Puerto Rico, Andrew W. Mellon foundation, and Smithsonian Tropical research Institute gave additional support. Data from the Pasoh Forest Reserve (Manokaran et al., 2004) were provided by the Forest Research Institute Malaysia and research was supported by the Negeri Sembilan State Forestry Department. The Yasuní forest dynamics plot was supported by the Pontifical Catholic University of Ecuador (Donaciones del Impuesto a la Renta from the government of Ecuador and PUCE grants L13251, M13373 in recent years). The Yasuní Scientific Station offered logistic support and the continuous research is endorsed by the Ministerio de Ambiente del Ecuador through several research permits (permits: No. 002-015-IC-FLO-PNY-DPAO, No. 025-2016-IC-FAU-FLODPAO-PNY, and No. 007-2018-IC-PNY-DPAO/AVS to Renato Valencia). Open Access funding enabled and organized by Projekt DEAL.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

DATA AVAILABILITY STATEMENT

Data are available by plot as follows. For the BCI plot: census data of trees dbh ≥ 1 cm (Condit et al., 2019) available in Dryad at <https://doi.org/10.15146/5xcp-0d46>; census data for seedling >20 -cm tall (Comita et al., 2023) available in a data paper published in Ecology at <https://doi.org/10.1002/ecy.4140>; species-level mean growth and survival rates of seedlings <20 tall (Kambach et al., 2024) available in iDiv, the repository of the German Centre for Integrative Biodiversity Research Halle-Jena-Leipzig, at <https://doi.org/10.25829/idiv.3565-ws9g97>; seed trap data (Wright & Calderon, 2015) available in the Smithsonian Tropical Research Institute repository at <https://doi.org/10.5479/si.data.201511251137>; trait data (Wright et al., 2016) available in Figshare at <https://doi.org/10.6084/m9.figshare.3550359.v1>. For the Fushan plot: species-level mean growth and survival rates of trees dbh ≥ 1 cm, species-level growth and survival rates of seedlings <50 -cm tall, and species-level mean annual seed numbers (Kambach et al., 2024) are available in iDiv at <https://doi.org/10.25829/idiv.3565-ws9g97>; trait data from Iida et al. (2014) at

<https://doi.org/10.1111/1365-2745.12221>. For the Luquillo plot: census data of trees dbh > 1 cm (Zimmerman, 2023) available in the EDI data portal at <https://doi.org/10.6073/pasta/7d937e27dfd99308362049d6c4495deb>; census data for seedling >20-cm tall (Zimmerman, 2018) available in the EDI data portal at <https://doi.org/10.6073/pasta/45e4817e74b51b9533b1bd4115415569>; seed trap data (Zimmerman & International Institute of Tropical Forestry, USDA Forest Service, 2023) available in the EDI data portal at <https://doi.org/10.6073/pasta/c0fb7433617bcd03dfd82bc1fad28cfb>; trait data (Swenson & Umana, 2015) available in Dryad at <https://doi.org/10.5061/dryad.j2r53>. For the Pasoh plot: data on growth and survival rates of trees dbh > 1 cm was collected in collaboration with the Forest Research Institute Malaysia (FRIM) and can only be obtained upon request (see original data request form in Appendix S1: Section S1, which was sent to prcsecretariat@frim.gov.my); species-level growth and survival rates of seedlings <50-cm tall (Kambach et al., 2024) available in the iDiv data repository at <https://doi.org/10.25829/ivdiv.3565-ws9g97>; and trait data (Wright & Yao, 2024) available in Dryad at <https://doi.org/10.5061/dryad.q2bvq83v7>. For the Yasuní plot: up to 200 growth observations and up to 1000 survival observations per species and size class dbh > 1 cm (Kambach et al., 2024) are available in the iDiv data repository at <https://doi.org/10.25829/ivdiv.3565-ws9g97>; growth and survival rates of seedlings <50 cm tall (Metz, Zambrano, et al., 2023) are available on the EDI data portal at <https://doi.org/10.6073/pasta/2cb969b626c3e276770a4fdc8bb3e375>; seed trap data (Garwood et al., 2023) are available on the EDI data portal at <https://doi.org/10.6073/pasta/5e6cb3d7ff741fd9d21965c4a904bc1f>; wood density data (Wright, Alvia, et al., 2024) are available in Figshare at <https://doi.org/10.25573/data.25639560.v2>; seed mass data (Garwood et al., 2024) are available on the EDI data portal at <https://doi.org/10.6073/pasta/95e4095ea61fb3cdc3b29b0cc10fa72e>; and specific leaf area data (Kraft & Ackerly, 2010) are available in Figshare at <https://doi.org/10.6084/m9.figshare.c.3292639>.

ORCID

Stephan Kambach  <https://orcid.org/0000-0003-3585-5837>

Liza S. Comita  <https://orcid.org/0000-0002-9169-1331>

Richard Condit  <https://orcid.org/0000-0003-4191-1495>

S. Joseph Wright  <https://orcid.org/0000-0003-4260-5676>

Margaret R. Metz  <https://orcid.org/0000-0002-4221-7318>

Simon A. Queenborough  <https://orcid.org/0000-0002-2468-0958>

REFERENCES

- Adler, P. B., R. Salguero-Gómez, A. Compagnoni, J. S. Hsu, J. Ray-Mukherjee, C. Mbeau-Ache, and M. Franco. 2014. "Functional Traits Explain Variation in Plant Life History Strategies." *Proceedings of the National Academy of Sciences of the United States of America* 111(2): 740–45.
- Anderson-Teixeira, K. J., S. J. Davies, A. C. Bennett, E. B. Gonzalez-Akre, H. C. Muller-Landau, S. J. Wright, K. A. Salim, et al. 2015. "CTFS-ForestGEO: A Worldwide Network Monitoring Forests in an Era of Global Change." *Global Change Biology* 21(2): 528–549.
- Bakewell, A. T., K. E. Davis, R. P. Freckleton, N. J. B. Isaac, and P. J. Mayhew. 2020. "Comparing Life Histories across Taxonomic Groups in Multiple Dimensions: How Mammal-Like Are Insects?" *The American Naturalist* 195(1): 70–81.
- Beukhof, E., R. Frelat, L. Pecuchet, A. Maureaud, T. S. Dencker, J. Sólmundsson, A. Punzón, et al. 2019. "Marine Fish Traits Follow Fast-Slow Continuum across Oceans." *Scientific Reports* 9(1): 17878.
- Blackburn, T. M. 1991. "Evidence for a 'Fast-Slow' Continuum of Life-History Traits among Parasitoid Hymenoptera." *Functional Ecology* 5(1): 65–74.
- Bogdziewicz, M., M.-C. A. Acuña, R. Andrus, D. Ascoli, Y. Bergeron, D. Brveiller, T. Boivin, et al. 2023. "Linking Seed Size and Number to Trait Syndromes in Trees." *Global Ecology and Biogeography* 32(5): 683–694.
- Bohman, S., and S. Pacala. 2012. "A Forest Structure Model that Determines Crown Layers and Partitions Growth and Mortality Rates for Landscape-Scale Applications of Tropical Forests." *Journal of Ecology* 100(2): 508–518.
- Browne, L., L. Markesteijn, E. Manzané-Pinzón, S. J. Wright, R. Bagchi, B. M. J. Engelbrecht, F. A. Jones, and L. S. Comita. 2023. "Widespread Variation in Functional Trait–Vital Rate Relationships in Tropical Tree Seedlings across a Precipitation and Soil Phosphorus Gradient." *Functional Ecology* 37: 248–260.
- Camargo, A. 2022. "PCAtest: Testing the Statistical Significance of Principal Component Analysis in R." *PeerJ* 10: e12967.
- Chang-Yang, C.-H., L. Chia-Ling, I. F. Sun, and C.-F. Hsieh. 2013. "Long-Term Seedling Dynamics of Tree Species in a Subtropical Rain Forest, Taiwan." *Taiwania* 58(1): 35–43.
- Chang-Yang, C.-H., P.-H. Chiang, S. J. Wright, C.-F. Hsieh, and I.-F. Sun. 2024. "Proximate Cues of Flowering in a Subtropical Rain Forest." *Biotropica* 56: 78–89.
- Chave, J., D. Coomes, S. Jansen, S. L. Lewis, N. G. Swenson, and A. E. Zanne. 2009. "Towards a Worldwide Wood Economics Spectrum." *Ecology Letters* 12: 351–366.
- Chen, Y.-Y. 2007. "Reproductive Phenology in a Lowland Dipterocarp Forest and Its Consequences for Seedling Recruitment." PhD thesis. Taipei, Taiwan: National Taiwan.
- Comita, L. S., S. Aguilar, S. P. Hubbell, and R. Pérez. 2023. "Long-Term Seedling and Small Sapling Census Data from the Barro Colorado Island 50 ha Forest Dynamics Plot, Panama." *Ecology* 104: e4140. <https://doi.org/10.1002/ecy.4140>.
- Comita, L. S., S. Aguilar, R. Pérez, S. Lao, and S. P. Hubbell. 2007. "Patterns of Woody Plant Species Abundance and Diversity in the Seedling Layer of a Tropical Forest." *Journal of Vegetation Science* 18(2): 163–174.

- Condit, R. 2022. "Expected Adult Lifespan in Tropical Trees: Long-Term Matrix Demography in a Large Plot." *Forest Ecosystems* 9: 100053.
- Condit, R., R. Pérez, S. Aguilar, S. Lao, R. Foster, and S. Hubbell. 2019. "Complete Data from the Barro Colorado 50-ha Plot: 423617 Trees, 35 Years." Dryad. <https://doi.org/10.15146/5xcp-0d46>.
- Condit, R., R. Pérez, S. Lao, S. Aguilar, and S. P. Hubbell. 2017. "Demographic Trends and Climate over 35 Years in the Barro Colorado 50 ha Plot." *Forest Ecosystems* 4(1): 17.
- Condit, R. S. 1998. *Tropical Forest Census Plots: Methods and Results from Barro Colorado Island, Panama, and a Comparison with Other Plots*. Berlin, New York: Springer.
- Condit, R. S., P. Ashton, S. Bunyavejchewin, H. S. Dattaraja, S. Davies, S. Esufali, C. Ewango, et al. 2006. "The Importance of Demographic Niches to Tree Diversity." *Science* 313(5783): 98–101.
- Cornelissen, J. H. C., S. Lavorel, E. Garnier, S. Díaz, N. Buchmann, D. E. Gurvich, P. B. Reich, et al. 2003. "A Handbook of Protocols for Standardised and Easy Measurement of Plant Functional Traits Worldwide." *Australian Journal of Botany* 51(4): 335–380.
- Davies, S. J., I. Abiem, K. A. Salim, S. Aguilar, D. Allen, A. Alonso, K. Anderson-Teixeira, et al. 2021. "ForestGEO: Understanding Forest Diversity and Dynamics through a Global Observatory Network." *Biological Conservation* 253: 108907.
- Daws, M. I., N. C. Garwood, and H. W. Pritchard. 2005. "Traits of Recalcitrant Seeds in a Semi-Deciduous Tropical Forest in Panama: Some Ecological Implications." *Functional Ecology* 19(5): 874–885.
- Dayrell, R. L. C., A. J. Arruda, S. Pierce, D. Negreiros, P. B. Meyer, H. Lambers, and F. A. O. Silveira. 2018. "Ontogenetic Shifts in Plant Ecological Strategies." *Functional Ecology* 32(12): 2730–41.
- de Kamimura, V. A., P. P. de Loiola, C. P. Carmona, M. A. Assis, C. A. Joly, F. A. M. Santos, S. A. Vieira, et al. 2023. "Trait Interactions Effects on Tropical Tree Demography Depend on the Environmental Context." *Perspectives in Plant Ecology, Evolution and Systematics* 59: 125732.
- Díaz, S., J. Kattge, J. H. C. Cornelissen, I. J. Wright, S. Lavorel, S. Dray, B. Reu, et al. 2016. "The Global Spectrum of Plant Form and Function." *Nature* 529(7585): 167–171.
- Fortunel, C., R. Valencia, S. J. Wright, N. C. Garwood, and N. J. B. Kraft. 2016. "Functional Trait Differences Influence Neighbourhood Interactions in a Hyperdiverse Amazonian Forest." *Ecology Letters* 19(9): 1062–70.
- Garwood, N. C., S. J. Wright, and M. R. Metz. 2024. "Seed Mass of Species from Yasuni National Forest, Ecuador, 2000–2014 ver 1." Environmental Data Initiative. <https://doi.org/10.6073/pasta/95e4095ea61fb3cdc3b29b0cc10fa72e>.
- Garwood, N. C., S. J. Wright, R. Valencia, and M. R. Metz. 2023. "Rainforest Phenology: Flower, Fruit and Seed Production from Biweekly Collections of 200 Traps in the Yasuni Forest Dynamics Plot, Ecuador, 2000–2018." Environmental Data Initiative. <https://doi.org/10.6073/pasta/5e6cb3d7ff741fd9d21965c4a904bc1f>.
- Gaiimo, S. 2014. "Evolution of Aging through Reduced Demographic Stochasticity – An Extension of the Pleiotropy Theory to Finite Populations." *Ecology and Evolution* 4(2): 167–173.
- Gibert, A., E. F. Gray, M. Westoby, I. J. Wright, and D. S. Falster. 2016. "On the Link between Functional Traits and Growth Rate: Meta-Analysis Shows Effects Change with Plant Size, as Predicted." *Journal of Ecology* 104(5): 1488–1503.
- Gilbert, B., S. J. Wright, H. C. Muller-Landau, K. Kitajima, and A. Hernández. 2006. "Life History Trade-Offs in Tropical Trees and Lianas." *Ecology* 87(5): 1281–88.
- He, Z., S. Webster, and S. Y. He. 2022. "Growth-Defense Trade-Offs in Plants." *Current Biology* 32(12): R634–R639.
- Hogan, J. A., J. K. Zimmerman, J. Thompson, M. Uriarte, N. G. Swenson, R. S. Condit, S. Hubbell, et al. 2018. "The Frequency of Cyclonic Wind Storms Shapes Tropical Forest Dynamism and Functional Trait Dispersion." *Forests* 9(7): 404.
- Hubbell, S. P., R. B. Foster, S. T. O'Brien, K. E. Harms, R. S. Condit, B. Wechsler, S. J. Wright, and S. Loo de Lao. 1999. "Light-Gap Disturbances, Recruitment Limitation, and Tree Diversity in a Neotropical Forest." *Science* 283: 554–57.
- Ickes, K., and S. C. Thomas. 2003. "Native, Wild Pigs (*Sus scrofa*) at Pasoh and Their Impacts on the Plant Community." In *Pasoh*, edited by T. Okuda, N. Manokaran, Y. Matsumoto, K. Niiyama, S. C. Thomas, and P. S. Ashton, 507–520. Tokyo: Springer Japan.
- Iida, Y., T. S. Kohyama, N. G. Swenson, S.-H. Su, C.-T. Chen, J.-M. Chiang, and I.-F. Sun. 2014. "Linking Functional Traits and Demographic Rates in a Subtropical Tree Community: The Importance of Size Dependency." *Journal of Ecology* 102: 641–650.
- John, J. A., and N. R. Draper. 1980. "An Alternative Family of Transformations." *Applied Statistics* 29(2): 190–97.
- Josse, J., and F. Husson. 2016. "missMDA: A Package for Handling Missing Values in Multivariate Data Analysis." *Journal of Statistical Software* 70(1): 1–31.
- Jucker, T., F. J. Fischer, J. Chave, D. A. Coomes, J. Caspersen, A. Ali, G. J. Loubota Panzou, et al. 2022. "Tallo: A Global Tree Allometry and Crown Architecture Database." *Global Change Biology* 28: 5254–68.
- Kambach, S., H. Bruehlheide, L. S. Comita, R. Condit, S. J. Wright, S. Aguilar, C.-H. Chang-Yang, et al. 2024. "Supplementary Data on Demographic Rates and Traits for Woody Plant Species at the Forest Dynamics Plots at Barro Colorado Island, Fushan, Pasoh and Yasuni (Version 1.0) [Dataset]." German Centre for Integrative Biodiversity Research. <https://doi.org/10.25829/idiv.3565-ws9g97>.
- Kambach, S., R. S. Condit, S. Aguilar, H. Bruehlheide, S. Bunyavejchewin, C.-H. Chang-Yang, Y.-Y. Chen, et al. 2022. "Consistency of Demographic Trade-Offs across 13 (Sub) Tropical Forests." *Journal of Ecology* 110(7): 1485–96.
- King, D. A., S. J. Wright, and J. H. Connell. 2006. "The Contribution of Interspecific Variation in Maximum Tree Height to Tropical and Temperate Diversity." *Journal of Tropical Ecology* 22: 11–24.
- Kohyama, T. S. 1993. "Size-Structured Tree Populations in Gap-Dynamic Forest-the Forest Architecture Hypothesis for the Stable Coexistence of Species." *Journal of Ecology* 81: 131.
- Kraft, N. J. B., and D. D. Ackerly. 2010. "Functional Trait and Phylogenetic Tests of Community Assembly across Spatial Scales in an Amazonian Forest." *Ecological Monographs* 80: 401–422.
- Kraft, N. J. B., and D. D. Ackerly. 2016. "Functional Trait and Phylogenetic Tests of Community Assembly across Spatial Scales in an Amazonian Forest." Wiley [Dataset Collection]. <https://doi.org/10.6084/m9.figshare.c.3292639.v1>.

- Kraft, N. J. B., R. Valencia, and D. D. Ackerly. 2008. "Functional Traits and Niche-Based Tree Community Assembly in an Amazonian Forest." *Science* 322: 580–82.
- Kunstler, G., D. A. Coomes, and C. D. Canham. 2009. "Size-Dependence of Growth and Mortality Influence the Shade Tolerance of Trees in a Lowland Temperate Rain Forest." *Journal of Ecology* 97(4): 685–695.
- Lande, R. 1982. "A Quantitative Genetic Theory of Life History Evolution." *Ecology* 63(3): 607–615.
- Lasky, J. R., B. Bachelot, R. Muscarella, N. Schwartz, J. Forero-Montaña, C. J. Nych, N. G. Swenson, J. Thompson, J. K. Zimmerman, and M. Uriarte. 2015. "Ontogenetic Shifts in Trait-Mediated Mechanisms of Plant Community Assembly." *Ecology* 96(8): 2157–69.
- Lasky, J. R., I.-F. Sun, S.-H. Su, Z.-S. Chen, and T. H. Keitt. 2013. "Trait-Mediated Effects of Environmental Filtering on Tree Community Dynamics." *Journal of Ecology* 101: 722–733.
- Legendre, P., and L. Legendre. 2012. *Numerical ecology*, 3rd ed. Amsterdam, Boston, MA: Elsevier.
- Lind, E. M., E. Borer, E. Seabloom, P. Adler, J. D. Bakker, D. M. Blumenthal, M. Crawley, et al. 2013. "Life-History Constraints in Grassland Plant Species: A Growth-Defence Trade-Off Is the Norm." *Ecology Letters* 16(4): 513–521.
- Luskin, M. S., J. S. Brashares, K. Ickes, I.-F. Sun, C. Fletcher, S. J. Wright, and M. D. Potts. 2017. "Cross-Boundary Subsidy Cascades from Oil Palm Degrade Distant Tropical Forests." *Nature Communications* 8: 2231.
- Manokaran, N., E. S. Quah, P. S. Ashton, and J. V. LaFrankie. 2004. "Pasoh Forest Dynamics Plot, Peninsular Malaysia." In *Tropical Forest Diversity and Dynamism: Findings from a Large-Scale Plot Network*, edited by E. C. Losos and E. G. Leigh, Jr., 585–598. Chicago, IL: University of Chicago Press.
- Metz, M. R., W. P. Sousa, and R. Valencia. 2010. "Widespread Density-Dependent Seedling Mortality Promotes Species Coexistence in a Highly Diverse Amazonian Rain Forest." *Ecology* 91(12): 3675–85.
- Metz, M. R., S. J. Wright, J. K. Zimmerman, A. Hernández, S. M. Smith, N. G. Swenson, M. N. Umaña, et al. 2023. "Functional Traits of Young Seedlings Predict Trade-Offs in Seedling Performance in Three Neotropical Forests." *bioRxiv* (preprint). <https://doi.org/10.1101/2023.01.14.523467>.
- Metz, M. R., M. Zambrano, R. Valencia, and N. C. Garwood. 2023. "Seedling Dynamics Demography Data, from the Yasuni Forest Dynamics Plot, Ecuador, 2002–2019." Environmental Data Initiative. <https://doi.org/10.6073/pasta/2cb969b626c3e276770a4fdc8bb3e375>.
- Muller-Landau, H. C. 2010. "The Tolerance-Fecundity Trade-Off and the Maintenance of Diversity in Seed Size." *Proceedings of the National Academy of Sciences of the United States of America* 107(9): 4242–47.
- Muscarella, R., M. Uriarte, J. Forero-Montaña, L. S. Comita, N. G. Swenson, J. Thompson, C. J. Nych, I. Jonckheere, and J. K. Zimmerman. 2013. "Life-History Trade-Offs during the Seed-to-Seedling Transition in a Subtropical Wet Forest Community." *Journal of Ecology* 101(1): 171–182.
- Needham, F. F., D. J. Johnson, K. J. Anderson-Teixeira, N. Bourg, S. Bunyavejchewin, N. Butt, M. Cao, et al. 2022. "Demographic Composition, Not Demographic Diversity, Predicts Biomass and Turnover across Temperate and Tropical Forests." *Global Change Biology* 28(9): 2895–2909.
- Nerlekar, A. N., and J. W. Veldman. 2020. "High Plant Diversity and Slow Assembly of Old-Growth Grasslands." *Proceedings of the National Academy of Sciences of the United States of America* 117(31): 18550–56.
- Ohse, B., A. Compagnoni, C. E. Farrior, S. M. McMahon, R. Salguero-Gómez, N. Rüger, and T. M. Knight. 2023. "Demographic Synthesis for Global Tree Species Conservation." *Trends in Ecology & Evolution* 38(6): 579–590.
- Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin, et al. 2020. "vegan: Community Ecology Package." <https://cran.r-project.org/web/packages/vegan/index.html>.
- Oli, M. K. 2004. "The Fast-Slow Continuum and Mammalian Life-History Patterns: An Empirical Evaluation." *Basic and Applied Ecology* 5(5): 449–463.
- Paine, C. E. T., L. Amissah, H. Auge, C. Baraloto, M. Baruffol, N. Bourland, H. Bruelheide, et al. 2015. "Globally, Functional Traits Are Weak Predictors of Juvenile Tree Growth, and We Do Not Know Why." *Journal of Ecology* 103(4): 978–989.
- Plummer, M., N. Best, K. Cowles, and K. Vines. 2006. "CODA: Convergence Diagnosis and Output Analysis for MCMC." *R News* 6: 7–11.
- Qiu, T., R. Andrus, M.-C. Aravena, D. Ascoli, Y. Bergeron, R. Berretti, D. Berveiller, et al. 2022. "Limits to Reproduction and Seed Size-Number Trade-Offs that Shape Forest Dominance and Future Recovery." *Nature Communications* 13: 2381.
- Qiu, T., M.-C. Aravena, R. Andrus, D. Ascoli, Y. Bergeron, R. Berretti, and M. Bogdziewicz. 2021. "Is There Tree Senescence? The Fecundity Evidence." *Proceedings of the National Academy of Sciences of the United States of America* 118(34): e2106130118.
- R Core Team. 2021. *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Reich, P. B. 2014. "The World-Wide 'Fast-Slow' Plant Economics Spectrum: A Traits Manifesto." *Journal of Ecology* 102(2): 275–301.
- Rüger, N., L. S. Comita, R. S. Condit, D. Purves, B. Rosenbaum, M. D. Visser, S. J. Wright, and C. Wirth. 2018. "Beyond the Fast-Slow Continuum: Demographic Dimensions Structuring a Tropical Tree Community." *Ecology Letters* 21(7): 1075–84.
- Rüger, N., R. S. Condit, D. H. Dent, S. J. DeWalt, S. P. Hubbell, J. W. Lichstein, O. R. Lopez, et al. 2020. "Demographic Trade-Offs Predict Tropical Forest Dynamics." *Science* 368(6487): 165–68.
- Russo, S. E., P. Brown, S. Tan, and S. J. Davies. 2008. "Interspecific Demographic Trade-Offs and Soil-Related Habitat Associations of Tree Species along Resource Gradients." *Journal of Ecology* 96(1): 192–203.
- Russo, S. E., S. M. McMahon, M. Detto, G. Ledder, S. J. Wright, R. S. Condit, S. J. Davies, et al. 2021. "The Interspecific Growth-Mortality Trade-Off Is Not a General Framework for Tropical Forest Community Structure." *Nature Ecology and Evolution* 5: 174–183.
- Salguero-Gómez, R., O. R. Jones, E. Jongejans, S. P. Blomberg, D. J. Hodgson, C. Mbeau-Ache, P. A. Zuidema, H. de Kroon, and Y. M. Buckley. 2016. "Fast-Slow Continuum and Reproductive Strategies Structure Plant Life-History Variation Worldwide." *Proceedings of the National Academy of Sciences of the United States of America* 113(1): 230–35.
- Stan Development Team. 2018. "RStan: The R Interface to Stan."
- Stearns, S. C. 1992. *The Evolution of Life Histories*. Oxford: Oxford University Press.

- Su, S. 2007. *Fushan Subtropical Forest Dynamics Plot: Tree Species Characteristics and Distribution Patterns*. Taipei: Taiwan Forestry Research Institute.
- Swenson, N. G., J. C. Stegen, S. J. Davies, D. L. Erickson, J. Forero-Montaña, A. H. Hurlbert, W. J. Kress, et al. 2012. "Temporal Turnover in the Composition of Tropical Tree Communities: Functional Determinism and Phylogenetic Stochasticity." *Ecology* 93: 490–99.
- Swenson, N. G., and M. N. Umana. 2015. "Data From: Interspecific Functional Convergence and Divergence and Intraspecific Negative Density Dependence Underlie the Seed-to-Seedling Transition in Tropical Trees [Dataset]." Dryad. <https://doi.org/10.5061/dryad.j2r53>.
- Umaña, M. N., J. Forero-Montaña, R. Muscarella, C. J. Nytych, J. Thompson, M. Uriarte, J. Zimmerman, and N. G. Swenson. 2016. "Interspecific Functional Convergence and Divergence and Intraspecific Negative Density Dependence Underlie the Seed-to-Seedling Transition in Tropical Trees." *The American Naturalist* 187: 99–109.
- Umaña, M. N., J. Needham, J. Forero-Montaña, C. J. Nytych, N. G. Swenson, J. Thompson, M. Uriarte, and J. K. Zimmerman. 2023. "Demographic Trade-Offs and Functional Shifts in a Hurricane-Impacted Tropical Forest." *Annals of Botany* 131: 1051–60.
- Uriarte, M., J. S. Clark, J. K. Zimmerman, L. S. Comita, J. Forero-Montana, and J. Thompson. 2012. "Multidimensional Trade-Offs in Species Responses to Disturbance: Implications for Diversity in a Subtropical Forest." *Ecology* 93(1): 191–205.
- Villellas, J., and M. B. García. 2013. "The Role of the Tolerance-Fecundity Trade-Off in Maintaining Intraspecific Seed Trait Variation in a Widespread Dimorphic Herb." *Plant Biology* 15(5): 899–909.
- Visser, M. D., M. Bruijning, S. J. Wright, H. C. Muller-Landau, E. Jongejans, L. S. Comita, and H. de Kroon. 2016. "Functional Traits as Predictors of Vital Rates across the Life Cycle of Tropical Trees." *Functional Ecology* 30(2): 168–180.
- Wickham, H. 2009. *ggplot2. Elegant Graphics for Data Analysis*. New York: Springer-Verlag New York.
- Wijenayake, P. R., T. Masaki, M. Shibata, and Y. Kubota. 2023. "Does Life Form Affect Tree Species Assembly? A Demographic Study across the Life History of a Temperate Forest in Japan." *Ecosphere* 14(6): e4579.
- Worthy, S. J., and N. G. Swenson. 2019. "Functional Perspectives on Tropical Tree Demography and Forest Dynamics." *Ecological Processes* 8(1): S165.
- Wright, J., and O. Calderon. 2015. "[Dataset:] Estimates of Seed Production Per Reproductive Basal Area Per Year for 38 Barro Colorado Island Tree Species for the Period 1993–2012." Smithsonian Tropical Research Institute. Dataset. <https://doi.org/10.5479/si.data.201511251137>.
- Wright, J. S., O. Calderon, and A. Hernandez. 2015. "Estimates of Mean Seed-to-Seedling Transition Probabilities for 68 Barro Colorado Island Tree Species for the Period 1993–2012."
- Wright, J. S., O. Calderón, A. Hernández, and H. C. Muller-Landau. 2024. "Flower and Seed Production, Seedling Dynamics, and Tree Life Cycles." In *The First 100 Years of Research on Barro Colorado Island: Plant and Ecosystem Science*, edited by H. C. Muller-Landau and S. J. Wright. Washington, DC: Smithsonian Institution Scholarly Press.
- Wright, J. S., K. Kitajima, N. J. B. Kraft, P. B. Reich, I. J. Wright, D. E. Bunker, R. S. Condit, et al. 2010. "Functional Traits and the Growth-Mortality Trade-Off in Tropical Trees." *Ecology* 91(12): 3664–74.
- Wright, J. S., K. Kitajima, N. J. B. Kraft, P. B. Reich, I. J. Wright, D. E. Bunker, R. S. Condit, et al. 2016. "Supplement 1. All Data Used in Analyses of Relationships among Functional Traits and Demographic Rates for Trees and Saplings from Barro Colorado Island, Panama." Wiley. Dataset. <https://doi.org/10.6084/m9.figshare.3550359.v1>.
- Wright, J. S., H. C. Muller-Landau, O. Calderón, and A. Hernández. 2005. "Annual and Spatial Variation in Seedfall and Seedling Recruitment in a Neotropical Forest." *Ecology* 86(4): 848–860.
- Wright, J. S., and T. L. Yao. 2024. "Plant Functional Traits of the 100 Most Abundant Species in the Pasoh Forest Reserve 50-ha Forest Dynamics Plot, Malaysia." Dryad. <https://doi.org/10.5061/dryad.q2bvq83v7>.
- Wright, S. J., P. Alvia, M. Zambrano, A. Perez, R. L. Valencia, and C. Altamirano. 2024. "Wood Density and Bark Thickness for 613 Tree Species from the 50-ha Forest Dynamics Plot at Yasuni, Ecuador." Smithsonian Tropical Research Institute. Dataset. <https://doi.org/10.25573/data.25639560.v2>.
- Zambrano, J., W. F. Fagan, S. J. Worthy, J. Thompson, M. Uriarte, J. K. Zimmerman, M. N. Umaña, and N. G. Swenson. 2019. "Tree Crown Overlap Improves Predictions of the Functional Neighbourhood Effects on Tree Survival and Growth." *Journal of Ecology* 107: 887–900.
- Zanne, A. E., G. Lopez-Gonzalez, D. A. Coomes, J. Ilic, S. Jansen, S. L. Lewis, R. B. Miller, et al. 2009. "Data From: Towards a Worldwide Wood Economics Spectrum." Dryad. <https://doi.org/10.5061/dryad.234>.
- Zimmerman, J. 2023. "Census of Species, Diameter and Location at the Luquillo Forest Dynamics Plot (LFDP), Puerto Rico." Environmental Data Initiative. <https://doi.org/10.6073/pasta/7d937e27dfd99308362049d6c4495deb>.
- Zimmerman, J., and International Institute of Tropical Forestry, USDA Forest Service. 2023. "Phenologies of the Tabonuco Forest Trees and Shrubs." Environmental Data Initiative. <https://doi.org/10.6073/pasta/c0fb7433617bcd03dfd82bc1fad28cfb>.
- Zimmerman, J. K. 2018. "LFDP Phenology Plot Seedlings-16 ha Plot. Version 5347491." Environmental Data Initiative. <https://doi.org/10.6073/pasta/45e4817e74b51b9533b1bd4115415569>.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Kambach, Stephan, Helge Bruelheide, Liza S. Comita, Richard Condit, S. Joseph Wright, Salomón Aguilar, Chia-Hao Chang-Yang, et al. 2025. "Putting Seedlings on the Map: Trade-Offs in Demographic Rates between Ontogenetic Size Classes in Five Tropical Forests." *Ecology* 106(1): e4527. <https://doi.org/10.1002/ecy.4527>